

# Sudan Agromet Dekadal Bulletin

Published by Sudan Meteorological Authority (SMA)

Vol 2, Issue 14

21-31 July 2004



## Highlights

- This dekad the ITCZ made remarkably turn southwards . The current position is about to 16°N, and it is remarkably south of average.
- Low or no significant rainfall occurred in most of North Darfur, North and West Kordofan and Kassala states. Areas with heavier rainfall were found mostly in the eastern border and parts of West and South Darfur.
- The rainfall totals relative to the long term average, remain below or on average. Central and North Sudan are mainly below average, but the bulk of the rain may come in August.
- In Greater Darfur and West Kordofan current evidence from the vegetation index and rainfall estimates indicates delays in the start of the growing season due to poor rains, but the situation is now improving – good rainfalls in August are important for recovery from this situation. In East Equatoria, Jonglei and Upper Nile, vegetation conditions are also markedly below average.

## Rainfall Analysis – Seasonal Progress

Rainfall in Sudan and its seasonal distribution is mostly the result of the northwards movement of moist air masses, source of the rainfall. The Intertropical Convergence Zone (ITCZ) marks the northernmost extent of these humid air masses, where they meet with drier and warmer air. The rains follow some distance south of this border between air masses, so that tracking this ITCZ through the season provides a quick evaluation of the seasonal movement of the rains




Fig 1a – Position of ITCZ over Africa in July Dek 3 2004 (red) compared to previous dekad (orange) and average position (black). Background is a rainfall map (Source : CPC-FEWS Net)




Fig 1b – Current latitude of the ITCZ position compared to the 15 year average. (Source : CPC-FEWS Net)

Note (fig 1a) how the ITCZ position marks the border between the (significantly) rainy and non-rainy rainfall areas. The way this position changes along the season can be described by the time series of its mean latitude (fig 1b). We can see :

- The ITCZ has been consistently north of the average for the past 3 dekads
- This dekad, it made a remarkable turn southwards (see Fig 1). The current position is about to 16°N.

## Rainfall Analysis - Dekadal Amounts and Frequency

10 day rainfall amounts produced by SAMIS at SMA/SEWS are based on a combination of METEOSAT satellite and synoptic gauge data. Rainfall climatology is similarly derived from a combination of historical data from the two sources.




Fig 2a – Rainfall amounts (mm) 21-31 July 2004




Fig 2b – Same but in relative terms : as % of long term average

In this dekad, the ITCZ moved southwards, the rainfall amounts were generally larger than previous dekad. Areas with heavier rainfall (over 80-100 mm) were found mostly along the eastern border of the country (Gedaref, Blue Nile and Sennar States) with point values of 128.1 mm reported at EL Damazin. Other areas of heavy rainfall are in parts of West and South Darfur with point values of 90.6 mm reported at Nyala.

In contrast, very low or no significant rainfall occurred in North Darfur, North and West Kordofan and Kassala states. Almost all of these regions registered less than 20mm.

In relative terms Eastern Sudan registered slightly more than the average rainfall, in contrast with Greater Kordofan and Southwestern Sudan where amounts were markedly below the average. Southeastern Sudan registered well above average rainfall, though absolute amounts were moderate.

## Rainfall Analysis - Cumulative Amounts

Cumulative amounts are obtained by summing the dekadal estimates starting from Dekad 1 of March until present.




Fig 3a – Cumulative rainfall (Mar Dek1 – Current Dek)




Fig 3b – Same but in relative terms : % of long term average


The cumulative rainfall amounts (Fig 3a) display the usual organisation in latitude bands (as the rainfall moves north following the ITCZ). Currently, values reach 700mm in southernmost Sudan and less than 50mm up to 16°N.

The situation relative to the average (Fig 3b) has not changed much from last dekad in many parts of Sudan. A belt of below average conditions running from Darfur across to Jonglei is noticeable. North Darfur, North and West Kordofan are the areas where below average conditions are more marked. In these areas it is important that rainfall situation during August improves.

Elsewhere, in the South and East of the country the situation is close to the average

## Vegetation Analysis

Vegetation information is based on the NDVI, a satellite index related to vegetation amount and vigour. NDVI data is sourced from the Africa Data Dissemination Service and processed at SMA.



*Fig 4a – NDVI 21-31 July 2004. Darker shades for denser vegetation, lightest shade for soil. (Source : ADDS).*

*Fig 4b – Same in relative terms : % of long term average (ADDS)*

The NDVI for this dekad (Fig 4a) continues the trend of vegetation advancing northwards following the progress of the rains. Currently, new season vegetation development is registered in the central states, up to the borders of Sennar, Gedaref, South and West Kordofan and West and South Darfur.

In Southern Sudan, vegetation conditions are mostly on average (Fig 4b) in broad accordance with the rainfall situation but declining vegetation conditions are noticeable in East Equatoria and Jonglei-Upper Nile borders.

In West Kordofan and Greater Darfur vegetation conditions are markedly below average. This reflects the same conditions in seasonal rainfall displayed in Fig 3b.

The indications from the vegetation index and rainfall estimates imply that early season vegetation development is somewhat delayed due to poor or irregular rains. However some improvements are already noticeable in Darfur but these are dependent on good rains during August.

## Acknowledgements

WFP Khartoum funded the installation of technical capacity and provision of training at SMA (Sudan Meteorological Authority) in the shape of SAMIS/SD – Satellite based Agro Meteorological Information System / Sudan. This is an operational system for the local production of satellite and station based rainfall and vegetation information for distribution to a wide range of local users. The system was prepared and installed by the TAMSAT group of the Department of Meteorology, Univ of Reading, UK. For information, please contact Dr. Rogério Bonifácio at : [tamsat@rdg.ac.uk](mailto:tamsat@rdg.ac.uk) or [rbonifacio@mail.telepac.pt](mailto:rbonifacio@mail.telepac.pt)

## Contacts

The SAMIS team includes :

Mr. Mousa Abdelbagi / Mrs. Hanan Awad Mohamed / Mrs. Badria Abdel Rahman

For further information, please contact

Dr. F. K. El-Sayem, Director General

Sudan Meteorological Authority

PO Box 574, Khartoum

Sudan

Tel. +249 11 778836/7 Fax. +249 11 771693

Contact : [su\\_samis@yahoo.com](mailto:su_samis@yahoo.com) / [ersad@sudanmail.net.sd](mailto:ersad@sudanmail.net.sd)